arXiv:1612.04195v1 [physics.gen-ph] 7 Dec 2016

Notes on Born-Infeld-type electrodynamics

S. I. Kruglov [1]

Department of Chemical and Physical Sciences, University of Toronto, 3359 Mississauga Road North, Mississauga, Ontario L5L 1C6, Canada

Abstract

We propose a new model of nonlinear electrodynamics with three parameters. Born-Infeld electrodynamics and exponential electrodynamics are particular cases of this model. The phenomenon of vacuum birefringence is studied. We show that there is not singularity of the electric field at the origin of point-like charged particles. The corrections to Coulomb's law at $r \to \infty$ are obtained. We calculate the total electrostatic energy of charges for different parameters of the model which is finite.

1 Introduction

For strong electromagnetic fields classical electrodynamics is modified due to self-interaction of photons [1]. Quantum electrodynamics with loop corrections becomes nonlinear electrodynamics (NLED) [2] and the effect of vacuum birefringence occurs in the presence of the external magnetic field. This phenomenon is now of experimental verification [3], [4], [5]. In Born-Infeld (BI) electrodynamics [6] there is no the effect of birefringence but in some NLED models the phenomenon of vacuum birefringence takes place [7]-[12]. The correspondence principle requires that in the limit of weak fields NLED has to be converted into Maxwell's electrodynamics. In classical electrodynamics problems of singularity of an electric field in the center of particles and the infinite electromagnetic energy of charges exist. The attractive feature of NLED [6], [10], [11], [13] is that there is an upper bound on the electric field at the origin of charges and the total electrostatic energy is finite. In this paper we propose a new model of NLED where the singularity is absent and the total electrostatic energy is finite.

The use units with $c = \hbar = 1$, $\varepsilon_0 = \mu_0 = 1$, and the metric signature is $\eta = \mathrm{diag}(-1, 1, 1, 1)$.

[1]E-mail: serguei.krouglov@utoronto.ca

2 A new model of nonlinear electrodynamics

Let us consider Born-Infeld-type electrodynamics with the Lagrangian density

$$\mathcal{L} = \frac{1}{\beta}\left[1 - \left(1 + \frac{\beta\mathcal{F}}{\sigma} - \frac{\beta\gamma\mathcal{G}^2}{2\sigma}\right)^{\sigma}\right], \tag{1}$$

where the parameters β and γ have the dimensions of (length)^4, and σ is the dimensionless parameter, $\mathcal{F} = (1/4)F_{\mu\nu}F^{\mu\nu} = (\mathbf{B}^2 - \mathbf{E}^2)/2$, $\mathcal{G} = (1/4)F_{\mu\nu}\tilde{F}^{\mu\nu} = \mathbf{E}\cdot\mathbf{B}$, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the field strength tensor, and $\tilde{F}^{\mu\nu} = (1/2)\epsilon^{\mu\nu\alpha\beta}F_{\alpha\beta}$ is the dual tensor. When $\sigma = 1/2$ the model (1) is converted to the model introduced in [8]. At $\beta = \gamma$ and $\sigma = 1/2$ we arrive at BI electrodynamics, The model described by the Lagrangian density (1) obeys the correspondence principle, i.e. at the weal field limit, $\beta\mathcal{F} \ll 1$, $\gamma\mathcal{G} \ll 1$, the NLED (1) is converted into Maxwell'l electrodynamics, $\mathcal{L}_M = -\mathcal{F}$. As a result, the nonlinearity of field equations is absent in the weak field limit. It should be noted that in the Born-Infeld-like model considered in [14] the correspondence principle does not hold for any parameter p introduced. At $\sigma \to \infty$ the model (1) is transformed into exponential electrodynamics

$$\mathcal{L}_{exp} = \lim_{\sigma\to\infty}\mathcal{L} = \frac{1}{\beta}\left[1 - \exp\left(\beta\mathcal{F} - \beta\gamma\mathcal{G}^2/2\right)\right]. \tag{2}$$

Other variants of exponential electrodynamics were proposed in [15], [12]. So, our model (1) allows us to consider simultaneously BI and exponential electrodynamics, as well as a new version of NLED.

2.1 Vacuum birefringence

Because in QED the effect of vacuum birefringence occurs [2], due to quantum corrections, it is of interest to study this effect in the model (1). The phenomenon of vacuum birefringence was measured in the BMV experiment [3] and in the PVLAS experiment [4]. In classical electrodynamics and in BI [6] electrodynamics vacuum birefringence is absent but in generalized BI electrodynamics with two parameters [8] the phenomenon of vacuum birefringence takes place. In the weak field limit $\beta\mathcal{F} \ll 1$, $\gamma\mathcal{G} \ll 1$, the Lagrangian density (1), making use of the Taylor series, becomes

$$\mathcal{L} = -\mathcal{F} + \frac{\beta(1-\sigma)}{2\sigma}\mathcal{F}^2 + \frac{\gamma}{2}\mathcal{G}^2$$

$$-\frac{\beta^2(\sigma-1)(\sigma-2)}{6\sigma^2}\mathcal{F}^3+\frac{\beta\gamma(\sigma-1)}{2\sigma}\mathcal{F}\mathcal{G}^2+\mathcal{O}\left((\beta\mathcal{F})^4,(\gamma\mathcal{G})^4\right). \qquad (3)$$

Up to $\mathcal{O}\left((\beta\mathcal{F})^2,(\gamma\mathcal{G})^2\right)$ the Lagrangian density (3) becomes the Heisenberg-Euler type [10]

$$\mathcal{L}_0=-\frac{1}{2}\left(\mathbf{B}^2-\mathbf{E}^2\right)+a\left(\mathbf{B}^2-\mathbf{E}^2\right)^2+b\left(\mathbf{E}\cdot\mathbf{B}\right)^2, \qquad (4)$$

where $a=\beta(1-\sigma)/8\sigma$, $b=\gamma/2$. The phenomenon of vacuum birefringence in the model with the Lagrangian density (4) was investigated in [10]. It was shown that the indices of refraction $n_\perp$, $n_\parallel$ for two polarizations, perpendicular and parallel to the external magnetic induction field $\bar{B}$, are

$$n_\perp=1+4a\bar{B}^2=1+\frac{\beta(1-\sigma)}{2\sigma}\bar{B}^2, \quad n_\parallel=1+b\bar{B}^2=1+\frac{\gamma}{2}\bar{B}^2. \qquad (5)$$

The phase velocities become $v_\perp=1/n_\perp$, $v_\parallel=1/n_\parallel$ and the phenomenon of vacuum birefringence holds when $n_\perp\neq n_\parallel$. According to the Cotton-Mouton (CM) effect [5] the $\triangle n_{CM}$ reads

$$\triangle n_{CM}=n_\parallel-n_\perp=k_{CM}\bar{B}^2. \qquad (6)$$

With the aid of Eqs. (5) and (6) one finds $k_{CM}=\gamma/2-\beta(1-\sigma)/2\sigma$. From the BMV and PVLAS experiments

$$k_{CM}=(5.1\pm6.2)\times10^{-21}\mathrm{T}^{-2} \qquad \text{(BMV)},$$
$$k_{CM}=(4\pm20)\times10^{-23}\mathrm{T}^{-2} \qquad \text{(PVLAS)}, \qquad (7)$$

we obtain the bound on our model parameters $\gamma/2-\beta(1-\sigma)/2\sigma\leq(4\pm20)\times10^{-23}\mathrm{T}^{-2}$. The effect of vacuum birefringence is absent if $k_{CM}=0$ ($\gamma=\beta(1-\sigma)/\sigma$). In BI electrodynamics $\gamma=\beta$, $\sigma=1/2$ and we obtain the known result that $k_{CM}=0$. In exponential electrodynamics, with the Lagrange density (2), when $\sigma\to\infty$ the phenomenon of vacuum birefringence disappears if $\beta=-\gamma$. The bound on CM coefficient in QED is given by $k_{CM}\leq4.0\times10^{-24}\mathrm{T}^{-2}$ [3].

3 Field equations

Making use of Euler-Lagrange equations we obtain equations of motion

$$\partial_\mu\left(\mathcal{L}_\mathcal{F}F^{\mu\nu}+\mathcal{L}_\mathcal{G}\tilde{F}^{\mu\nu}\right)=0, \qquad (8)$$

where $\mathcal{L}_{\mathcal{F}} = \partial\mathcal{L}/\partial\mathcal{F}$, $\mathcal{L}_{\mathcal{G}} = \partial\mathcal{L}/\partial\mathcal{G}$, and $\tilde{F}^{\mu\nu} = (1/2)\epsilon^{\mu\nu\alpha\beta}F_{\alpha\beta}$ is the dual tensor. By virtue of Eqs. (1) and (8) one finds field equations

$$\partial_\mu \left[\Pi^{\sigma-1}\left(F^{\mu\nu} - \gamma\mathcal{G}\tilde{F}^{\mu\nu}\right)\right] = 0, \qquad \Pi = 1 + \frac{\beta\mathcal{F}}{\sigma} - \frac{\beta\gamma\mathcal{G}^2}{2\sigma}. \tag{9}$$

The electric displacement field is given by $\mathbf{D} = \partial\mathcal{L}/\partial\mathbf{E}$,

$$\mathbf{D} = \Pi^{\sigma-1}\left(\mathbf{E} + \gamma\mathcal{G}\mathbf{B}\right). \tag{10}$$

The magnetic field obtained from the relation $\mathbf{H} = -\partial\mathcal{L}/\partial\mathbf{B}$ is

$$\mathbf{H} = \Pi^{\sigma-1}\left(\mathbf{B} - \gamma\mathcal{G}\mathbf{E}\right). \tag{11}$$

We can decompose Eqs. (10) and (11) as [16]

$$\mathbf{D} = \varepsilon\mathbf{E} + \nu\mathbf{B}, \qquad \mathbf{H} = \mu^{-1}\mathbf{B} - \nu\mathbf{E}. \tag{12}$$

From Eqs. (10), (11) and (12) one finds

$$\varepsilon = \Pi^{\sigma-1}, \qquad \mu^{-1} = \varepsilon, \qquad \nu = \gamma\mathcal{G}\Pi^{\sigma-1}. \tag{13}$$

With the help of Eqs. (10) and (11), field equations (9) may be written in the form of nonlinear Maxwell's equations

$$\nabla\cdot\mathbf{D} = 0, \qquad \frac{\partial\mathbf{D}}{\partial t} - \nabla\times\mathbf{H} = 0. \tag{14}$$

As ε, μ^{-1}, and ν depend on electromagnetic fields equations (14) are nonlinear Maxwell's equations. The Bianchi identity gives the second pair of nonlinear Maxwell's equations

$$\nabla\cdot\mathbf{B} = 0, \qquad \frac{\partial\mathbf{B}}{\partial t} + \nabla\times\mathbf{E} = 0. \tag{15}$$

Making use of Eqs. (12) and (13) we obtain the equality

$$\mathbf{D}\cdot\mathbf{H} = (\varepsilon^2 - \nu^2)\mathbf{E}\cdot\mathbf{B} + 2\varepsilon\nu\mathcal{F}. \tag{16}$$

Because $\mathbf{D}\cdot\mathbf{H} \neq \mathbf{E}\cdot\mathbf{B}$ [17] the dual symmetry is broken in our model as well as in generalized BI electrodynamics [8] and in QED with quantum corrections. In classical electrodynamics and in BI electrodynamics the dual symmetry occurs.

3.1 The electric field of point-like charged particles

Adding the source in Eq. (14) for the point-like charged particle with the electric charge Q, in Gaussian units, we obtain the equation

$$\nabla \cdot \mathbf{D} = 4\pi Q\delta(\mathbf{r}). \tag{17}$$

The solution to Eq. (17) is

$$\mathbf{D} = \frac{Q}{r^3}\mathbf{r}. \tag{18}$$

By virtue of Eq. (10) it becomes

$$\frac{E}{[1-\beta E^2/(2\sigma)]^{1-\sigma}} = \frac{Q}{r^2}. \tag{19}$$

We make a conclusion that if $\sigma < 1$ at $r \to 0$ the solution to Eq. (19) is given by

$$E(0) = \sqrt{\frac{2\sigma}{\beta}}. \tag{20}$$

At $\sigma = 1/2$ we come to known result that in BI electrodynamics the maximum field in the origin of charged particle is $E(0) = \sqrt{1/\beta}$. For the case of $\sigma > 1$, as in exponential electrodynamics ($\sigma \to \infty$), there is a singularity in the center of particles. At $\sigma < 1$, there is no singularity of the electric field strength at the origin of point-like charged particles and Eq. (20) gives the maximum of the electric field in the center of charged particles. Let us introduce unitless variables

$$x = \frac{(2\sigma)^{1/4}}{\sqrt{Q}\beta^{1/4}}r, \qquad y = \sqrt{\frac{\beta}{2\sigma}}E. \tag{21}$$

Using Eqs. (21) equation (19) takes the form

$$\frac{y}{(1-y^2)^{1-\sigma}} = \frac{1}{x^2}. \tag{22}$$

The plot of the function $y(x)$ for $\sigma = 0.1$, $\sigma = 0.5$ and $\sigma = 0.8$ is represented by Fig. 1. The Taylor series of the function $y(x)$ at $x \to \infty$ ($r \to \infty$) reads

$$y = \frac{1}{x^2} - \frac{1-\sigma}{x^6} + \mathcal{O}(x^{-8}). \tag{23}$$



Figure 1: The plot of the function y vs. x.

From Eqs. (21) and (23) one can obtain the asymptotic value of the electric field at $r \to \infty$

$$E(r) = \frac{Q}{r^2} - \frac{(1-\sigma)\beta Q^3}{2\sigma r^6} + \mathcal{O}(r^{-8}). \tag{24}$$

Eq. (24) shows the correction to Coulomb's law at $r \to \infty$ for any value of the parameter σ. For BI electrodynamics, when $\sigma = 1/2$, we find the asymptotic of the electric field

$$E(r) = \frac{Q}{r^2} - \frac{\beta Q^3}{2r^6} + \mathcal{O}(r^{-8}) \qquad (\sigma = 1/2). \tag{25}$$

For exponential electrodynamics ($\sigma \to \infty$) one obtains from Eq. (24) the correction to Coulomb's law similar to BI electrodynamics but with the opposite sign, $E(r) = Q/r^2 + \beta Q^3/(2r^6) + \mathcal{O}(r^{-8})$. At $\beta = 0$ we come to Maxwell's electrodynamics and the Coulomb law $E(r) = Q/r^2$ is recovered. The electric field singularities are absent at the origin of charged particles at $\sigma < 1$.

4 Energy-momentum tensor and the energy of charged particles

The general expression for the symmetrical energy-momentum tensor is given by

$$T_{\mu\nu} = \left(\mathcal{L}_{\mathcal{F}} F_{\mu}^{\ \alpha} + \mathcal{L}_{\mathcal{G}} \tilde{F}_{\mu}^{\ \alpha}\right) F_{\nu\alpha} - g_{\mu\nu}\mathcal{L}. \tag{26}$$

From Eqs. (1) and (26) we find the symmetrical energy-momentum tensor

$$T_{\mu\nu} = -\Pi^{\sigma-1}\left(F_{\mu}^{\ \alpha} F_{\nu\alpha} - \gamma\mathcal{G}\tilde{F}_{\mu}^{\ \alpha} F_{\nu\alpha}\right) - g_{\mu\nu}\mathcal{L}. \tag{27}$$

For the case $\mathbf{B} = 0$ the electric energy density becomes

$$\rho_E = T_0^{\ 0} = \frac{E^2}{(1-\beta E^2/(2\sigma))^{1-\sigma}} + \frac{1}{\beta}\left[\left(1 - \frac{\beta E^2}{2\sigma}\right)^{\sigma} - 1\right]. \tag{28}$$

The total electrostatic energy of point-like charges is given by $\mathcal{E} = \int_0^\infty \rho_E r^2 dr$. Making use of Eqs. (21), (22) and (28), and changing variables, we obtain

$$\mathcal{E} = \frac{Q^{3/2}}{2^{7/4}\sigma^{3/4}\beta^{1/4}} \int_0^1 \left[1 - \frac{2\sigma y^2}{(1-y^2)^{1-\sigma}} - (1-y^2)^{\sigma}\right] \times \frac{(1-y^2)^{(1-3\sigma)/2}\left[(2\sigma-1)y^2 - 1\right] dy}{y^{5/2}}. \tag{29}$$

The normalized electrostatic energy $\mathcal{E}\beta^{1/4}/Q^{3/2}$ of point-like charges for different parameters σ, found from Eq. (29), is given in Table 1. Thus, the

Table 1: The normalized electrostatic energy

σ	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8
$\mathcal{E}\beta^{1/4}/Q^{3/2}$	0.788	0.947	1.061	1.154	1.250	1.311	1.382	1.451

total electrostatic energy of point-like particles is finite for $0 < \sigma < 1$.

5 Conclusion

We have proposed a new model of NLED with three parameters β, γ and σ and BI electrodynamics (at $\beta = \gamma$, $\sigma = 1/2$) and exponential electrodynamics (at $\sigma \to \infty$) are particular cases of this model. The correspondence principle takes place so that for weak fields the model is converted to Maxwell's electrodynamics. The phenomenon of vacuum birefringence occurs if $\gamma \neq \beta(1-\sigma)/\sigma$ otherwise the effect of birefringence disappears. We demonstrated that the dual symmetry is violated in our model. It is shown that there is no singularity of the electric field in the center of point-like charged particles and the maximum electric field strength is $E(0) = \sqrt{2\sigma}/\sqrt{\beta}$. The corrections to Coulomb's law at $r \to \infty$ are obtained which are in the order of $\mathcal{O}(r^{-6})$. We calculated the total electrostatic energy of charged particles for different values of the parameter σ, that is finite.

References

[1] J. D. Jackson, *Classical Electrodynamics, Second Ed.*, John Wiley and Sons, 1975.

[2] W. Heisenberg and H. Euler, Z. Physik, **98**, 714 (1936) (arXiv:physics/0605038).

[3] A. Cadene, P. Berceau, M. Fouche, R. Battesti and C. Rizzo, *Eur. Phys. J. D* **68**, 16 (2014) (arXiv:1302.5389).

[4] F. Della Valle, et al, *Phys. Rev. D* **90**, 092003 (2014).

[5] R. Battesti and C. Rizzo, *Rep. Prog. Phys.* **76**, 016401 (2013) (arXiv:1211.1933).

[6] M. Born and L. Infeld, Proc. Royal Soc. (London) A **144**, 425 (1934).

[7] S. I. Kruglov, Phys. Rev. D **75**, 117301 (2007).

[8] S. I. Kruglov, J. Phys. A **43**, 375402 (2010) (arXiv:0909.1032).

[9] S. I. Kruglov, Eur. Phys. J. C **75**, 88 (2015) (arXiv:1411.7741 [hep-th]).

[10] S. I. Kruglov, Ann. Phys. (Berlin) **527**, 397 (2015) (arXiv:1410.7633); Commun. Theor. Phys. **66**, 59 (2016) (arXiv:1511.03303 [hep-ph]).

[11] S. I. Kruglov, Ann. Phys. **353**, 299 (2015) (arXiv:1410.0351); Phys. Lett. A **379**, 623 (2015) (arXiv:1504.03535).

[12] S. I. Kruglov, Int. J. Mod. Phys. A **31**, 1650058 (2016) (arXiv:1607.03923 [gr-qc]); Europhys. Lett. **115**, 60006 (2016) (arXiv:1611.02963).

[13] D. M. Gitman and A. E. Shabad, Eur. Phys. J. C **74**, 3186 (2014).

[14] P. Gaete and J. Helayl-Neto, Eur. Phys. J. C **74**, 3182 (2014) (arXiv:1408.3363).

[15] S. H. Hendi, JHEP **03**, 065 (2012).

[16] F. W. Hehl and Yu. N. Obukhov, *Foundations of classical electrodynamics: Charge, flux, and metric* (Birkhäuser, Boston, 2003).

[17] G. W. Gibbons and D. Rasheed, *Nucl. Phys. B* **454** (1995) 185 (arXiv:hep-th/9506035).